

AC JOINT RECONSTRUCTION PROTOCOL

DR ELRASHIDY – TRI-VALLEY ORTHOPEDICS

Procedure: **Coracoclavicular Reconstruction with Semitendinosis Graft**

Frequency: 2 visits/week **Duration:** 4 months

Post-Op: Sling x 6 weeks, Cryocuff x 2 weeks, Dressing Change at 48 hours

Phase I (Weeks 0-2): Post-Operative

Precautions

- Sling or Immobilizer: At all times except with PT or when showering.
- No shoulder ROM
- No lifting, no body weight support with hands

Exercises

- Begin Pendulums as tolerated
- AROM wrist/elbow/digits
- Scapular “pinches”

Phase II (Weeks 3-6): Protection Phase

Precautions (Through Week 6)

- Sling or Immobilizer: At all times except with PT or when showering.
- No lifting (nothing greater than a coffee cup)

Exercises:

- Passive supine ER to neutral and extension to neutral
- Passive supine FF in scapular plane to 100°
- AROM wrist/elbow, Scapular “pinches”
- Pain free submaximal deltoid isometrics

Phase III (Weeks 6-10): Intermediate/Range of Motion Phase

- Sling/Immobilizer: **Discontinue at Week 6.**

Exercises:

- Passive & Active-Assisted FF in scapular plane - limit 140° (wand exercises, pulleys)
- Passive & Active assisted ER - No limits (go SLOW with ER)
- Manual scapular side-lying stabilization exercises, IR/ER submaximal
- Pain free isometrics and soft tissue modalities as needed

Advancement Criteria:

- FF to 160°, ER to 40°

AC JOINT RECONSTRUCTION

- Normal scapulohumeral rhythm
- Minimal pain and inflammation

Phase IV (Weeks 10-14): Early Strengthening

Exercises:

- AAROM for full FF and ER, AAROM for IR - no limits
- IR/ER/FF isotonic strengthening
- Scapular and latissimus strengthening
- Humeral head stabilization exercises
- Begin biceps strengthening
- Progress IR/ER to 90/90 position if required
- General upper extremity flexibility exercises

Advancement Criteria:

- Normal scapulohumeral rhythm and full upper extremity ROM
Isokinetic IR/ER strength 85% of uninvolved side
- Minimal pain and inflammation

Phase V (Weeks 14-18): Advance Strengthening/Return to Activity

Exercises:

- Continue full upper extremity strengthening program
- Continue upper extremity flexibility exercises
- Activity-specific plyometrics program
- Begin sport or activity related program
- Address trunk and lower extremity demands
- Begin throwing program if appropriate
 - Begin light ball toss @ 20-30ft. max, 60% velocity, work on wind-up mechanics, early & late cocking phase, acceleration & follow through
 - Throwers begin re-entry throwing program on level surface (see interval return to throwing program)
 - Continue strengthening and stretching programs
- Emphasize posterior capsule stretching

Discharge Criteria:

- Isokinetic IR/ER strength equal to uninvolved side
- Independent HEP
- Independent, pain-free sport or activity specific program

AC JOINT RECONSTRUCTION PROTOCOL


DR ELRASHIDY – TRI-VALLEY ORTHOPEDICS

EXERCISES

- 1) **Passive Supine Flexion/Forward Elevation:** Lie on your back. Grasp wrist with non-op hand and passively raise operative arm overhead. Aim to get to 90° by 3 weeks. In week 4, progress to 120°. Then full ROM after 6 weeks. Keep elbow bent and relaxed. Repeat 10 reps, 2-3 times/day



- 2) **Passive Pendulum Exercise:** Hold onto a chair back with non-op hand and bend forward. Let the operative arm hang down passively. Use body to passively swing arm: Forward, backward, side to side and in small circles. Repeat throughout the day as tolerated



Hany Elrashidy. MD
Sports Medicine, Arthroscopy
Department of Orthopedic Surgery
Tri-Valley Orthopedic Specialists
www.hanyelrashidymd.com