
SLAP REPAIR PROTOCOL

DR ELRASHIDY – TRI-VALLEY ORTHOPEDICS

- ◆ Primary goal of PT: Protect SLAP repair
- ◆ Pain is NOT a good gauge of progression, healing time constraints CRITICAL

PHASE I: Weeks 0-6 (Protected Motion Phase)

SLING FULL-TIME for 6 weeks

CRYOTHERAPY X 14 days

PROM @ Elbow & Forearm – NO isolated BICEPS contraction/lifting x 6 weeks

NO PULLEYS unless noted by MD (and then with elbow passively flexed to 60°)

No repetitive ER/IR (places too much stress at repair)

No Active Motion until 6 weeks post-op

No Strengthening or Isometrics until 12 weeks post-op

ROM LIMITS

- ◆ Sling for 6 weeks (unless otherwise noted by MD)
 - ◆ AROM of wrist & hand
 - ◆ Soft tissue modalities to axilla, pec major/minor for pain/swelling prn
 - ◆ Initiate pre-setting of scapula and scapular stabilizers for posture re-education
- Include scapular retraction/posture/shrugs/scapular clocks
- ◆ **Shoulder PROM** only: Focus on Passive Supine Forward Elevation in first 6 weeks as well as pendulums

Week 0-2 (Avoid EXTENSION & ABDUCTION)

- ◆ **FLEXION:** Up to 60°
- ◆ **ELEVATION:** Up to 60° in scapular plane
- ◆ **ER:** Up to 0° MAXIMUM (Peel-Back Mechanism)
- ◆ **IR:** Up to 45° maximum in scapular plane with elevation to 60°

Week 3-4 (Avoid EXTENSION & ABDUCTION WITH ER ROM)

- FLEXION:** Up to 60°
- ABDUCTION:** Up to 60° with UE IR (Peel-Back Mechanism)
- ER:** Up to 30° in scapular plane with elevation up to 60°. AVOID AROM ER
- IR:** Up to 60° in scapular plane with elevation up to 60°

Week 5-6 – Gradually improve PROM & Progress:

- FLEXION:** Up to 120°
- ABDUCTION:** Up to 60° with UE IR (Peel-Back Mechanism)
- ER:** Up to 50° in scapular plane with elevation up to 60°. AVOID AROM ER
- IR:** Up to 60° in scapular plane with elevation up to 60°

- ◆ Initiate light posterior capsular stretching (Horizontal adduction below 90° elevation)
- ◆ Continue to work on scapular pre-setting for posture re-education

SLAP REPAIR PROTOCOL

PHASE II: Weeks 7-12 (Moderate Protection Phase)

- ◆ D/C sling
- ◆ Allow use of operative extremity for light ADLs. **NO Lifting.**
- ◆ Gradually progress to AROM: Straight plane movements

Weeks 7-9:

- ◆ **FLEXION:** Up to 180°
- ◆ **ABDUCTION:** Up with UE ER to tolerance
- ◆ **ER:** Up to 90° at 90° abduction
- ◆ **IR:** Up to 70° at 90° abduction
- ◆ Begin extension A/PROM and begin Biceps AROM
- ◆ Begin light pain-free isometrics for shoulder musculature (ABD, ER/IR, Extension with UE @ side allowed.
- ◆ No biceps, forward flexion or elevation isometrics

Weeks 10-12:

- ◆ Progress ROM to functional demands of patient (i.e overhead athlete; combined movements)
- ◆ Progress ER to thrower's motion: ER up to 115° at 90° abduction

PHASE III: Weeks 13-20 (Minimal Protection Phase)

Weeks 13-16:

- ◆ Initiate Isotonic strengthening program with progressive strengthening of rotator cuff and scapular stabilizers
 - ◆ Begin Biceps sub-maximal Isometrics then progress to Isotonic strengthening of Biceps as well as supination exercises
 - ◆ Start light shoulder and UE strengthening (No < 5 lbs)
- Continue all stretching and maintain thrower's motion (ER)
- ◆ Initiate weights for RC strengthening as well as serratus, mid/low trap, bi/triceps
 - ◆ T-Band IR and ER @ side
 - ◆ Restricted sports activities (light swimming, half golf swings)
 - ◆ Independent shoulder stretches and light plyometric program

Weeks 17-20:

- ◆ Initiate interval throwing program on level ground

SLAP REPAIR PROTOCOL
DR ELRASHIDY – TRI-VALLEY ORTHOPEDICS

PHASE IV: Week 21-26 (Advanced Strengthening)

- ◆ Initiate T-Band ER/IR at 90 Abduction (slow/fast sets)
- ◆ Continue strengthening program to progress endurance
- ◆ Progress Interval throwing program and/or initiate Sports-specific drills (including long-toss)

Phase V: Months 6-9 (RETURN TO SPORT)

- ◆ Advanced interval throwing program from pitcher's mound
- ◆ Continue/advance sports-specific drills

Hany Elrashidy. MD
Sports Medicine, Arthroscopy
Department of Orthopedic Surgery
Tri-Valley Orthopedic Specialists
www.hanyelrashidymd.com


SLAP REPAIR PROTOCOL

EXERCISES: PHASE I


- 1) **Passive Supine Flexion/Forward Elevation:** Lie on your back. Grasp wrist with non-op hand and passively raise operative arm overhead. Aim to get to 90° by 3 weeks. In week 4, progress to 120°. Then full ROM after 6 weeks. Keep elbow bent and relaxed. Repeat 10 reps, 2-3 times/day



- 2) **Passive Pendulum Exercise:** Hold onto a chair back with non-op hand and bend forward. Let the operative arm hang down passively. Use body to passively swing arm: Forward, backward, side to side and in small circles. Repeat throughout the day as tolerated



- 3) **Passive Elbow Flexion/Extension:** While standing or sitting, use non-op hand to bend & straighten elbow. Repeat 10 reps, 2-3 times/day




SLAP REPAIR PROTOCOL

DR ELRASHIDY – TRI-VALLEY ORTHOPEDICS

EXERCISES: PHASE II

1) Active external rotation (NO Weight): Lie supine or on non-op side. With elbow bent to 90, rotate upper arm up and down (per guidelines). Repeat 5-10 reps, 2-3 times/day



2) Posterior capsular stretch: Gently reach across body. Use other hand to pull elbow further across and hold for 10-20 seconds, 1-2 times/day



3) Wall Climb Abduction: Stand with shoulders flush with doorway. Use thumb & fingers to climb sideways up wall as high as possible. Do 10-20 reps, 2-3 times/day



EXERCISES: PHASE III

1) Theraband ER and IR: Secure knotted band in a door and grasp with hand. Keeping elbow bent, and in at side, pull band towards stomach, followed by pulling out with hand. Do 10-20 reps, 1-2 times/day

